

Read this sheet in full before using, and keep for later reference.

Find more product info online

Stoppa

Search

Class 2
OTC drugs

Stoppa EX Antidiarrheal <Grapefruit Flavor>

Effective against sudden diarrhea and diarrhea accompanied by pain.

Dissolves in the mouth without water,

making it convenient to take while on the train or wherever you happen to be.

Relief with a single tablet taken without water

Contains no drowsiness-inducing ingredients

(However, users should do not drive or operate machinery)

For relief from diarrhea relief when out and about



Meetings or tests



When travelling



After eating or drinking too much

Excellent Effectiveness

The herbal medicine **scopolia extract** helps reduce abnormal intestinal contractions, slowing the movement of stool through the intestines. At the same time, **berberine tannate** helps reduce inflammation of the intestinal mucosa and kill diarrhea-causing bacteria to help treat diarrhea caused by food or water contamination.



Scopolia root

Dissolves Fast

Can be taken anywhere

The “**express**” tablets quickly dissolve in the mouth. Easy to swallow without water, the tablets can be taken whenever, wherever.

EXPRESS Tablet



(Artist's rendering)

Warnings

Do not

(Failure to comply with these instructions increases the likelihood of current symptoms worsening, side effects occurring, and other potentially serious risks)

1. **Do not take any of the following pharmaceuticals while using this product**

Gastrointestinal antispasmodic pain relievers; other gastrointestinal medicines that contain scopolia extract; motion sickness medications

2. **Do not drive or operate machinery while using this product**

(users may experience blurred vision, abnormal sensitivity to light or other similar symptoms)

3. **Individuals who are breastfeeding should avoid taking this product, or avoid breastfeeding after taking the product**

(ingredients may transfer to breast milk and cause and cause an elevated heartbeat in nursing infants)

Seek advice

1. Ask a doctor, pharmacist or registered seller before use if you are

- (1) Currently receiving treatment from a doctor
- (2) Have diarrhea accompanied by a fever, bloody stool or persistent mucous stool
- (3) Have sudden, severe diarrhea or have diarrhea accompanied by abdominal pain, abdominal swelling, nausea, etc. (Stopping diarrhea with this product may cause illness to worsen.)
- (4) Pregnant or suspect you may be pregnant
- (5) Elderly
- (6) Have ever shown allergic symptoms when taking a drug or medicine
- (7) Have the following symptom: Difficulty urinating
- (8) Have been diagnosed with any of the following conditions: Heart disease; glaucoma

2. The following symptoms may be side effects of the product. If you experience any of the following symptoms when using the product, stop use immediately and bring this sheet to a doctor, pharmacist or registered seller for consultation.

Affected area	Symptom
Skin	Rash or redness; itchiness
Mental state / Nervous system	Headache
Urinary organs	Difficulty urinating
Other	Flushed face; abnormal sensitivity to light

3. The following symptoms may occur after using the product. If these symptoms persist or worsen, stop use and bring this sheet to a doctor, pharmacist or registered seller for consultation.

Dry eyes; blurred vision

4. If symptoms do not improve after 5 to 6 days of use, stop use and bring this sheet to a doctor, pharmacist or registered seller for consultation.

Other warnings

May result in reduced breast milk flow.

Effects

Relieves diarrhea accompanied by abdominal pain; diarrhea; diarrhea due to indigestion; food and water poisoning; diarrhea with vomiting; loose bowels; loose stool.

Directions and Dosage

Take the following amount by chewing or dissolving the tablet in the mouth.

Age	Amount per 1 dose	Doses per day	Time between doses
Adults (15 years and up)	1 tablet	No more than 3	At least 4 hours
Children under 15 years	✗ Do not use		

Usage and Dosage Warnings

- (1) Use only as directed.
- (2) To remove the tablets from the package: As shown in the illustration on the right, press firmly with a fingertip on the protruding bubble in the clear plastic sheet that contains the tablet, pushing the tablet through the foil backing and out (swallowing the product without first removing it from the package can lead to injury, such as damage to the esophageal membrane).



Ingredients (per 1 tablets)

Active ingredient	Content	Purpose
1:2 trituration of scopolia extract (20 mg scopolia extract)	60 mg	Helps reduce abnormal intestinal contractions, slowing the movement of stool in the intestines. Highly effective for diarrhea accompanied by abdominal pain.
Berberine tannate	100 mg	Helps protect the intestinal mucosa and reduce inflammation while helping prevent excess water from accumulating in the intestines. Also helps suppress abnormal decomposition and fermentation within the intestines.

Inactive ingredients: D-mannitol, cellulose, crospovidone, cornstarch, gum Arabic, magnesium stearate, aspartame (an L-phenylalanine compound), menthol and flavoring.

Note about ingredients

Because this product contains herbal medicine (plant roots, bark, etc., with medicinal properties), individual tablets may vary somewhat in color or appear to contain black flecks, but these variations do not affect the product's efficacy.

Reason for the tablets' yellow color

The berberine tannate contained in Stoppa EX Antidiarrheal tablets causes the tablets to appear yellow. Tablets contain no added coloring.

Storage and Handling

- (1) Store out of direct sunlight in a cool, dry place.
- (2) Keep out of reach of children.
- (3) Do not transfer to another container or package (may lead to incorrect use or affect product quality).
- (4) Do not use after the use-by date indicated on the package.
- (5) Avoid damaging the foil backing on the packaging for tablets you are not planning to use immediately, especially when tearing the plastic sheets containing the tablets along perforated lines, as damage to the packaging can affect product quality.

Inquiries

Please direct inquiries to the store where you purchased the product or the service center below.

Lion Corporation Consumer Service Center [0120-813-752](tel:0120-813-752)

Hours available: 9:00–17:00 (excluding Saturdays, Sundays and holidays)

Lion Corporation

3-7, Honjo 1-chome, Sumida-ku, Tokyo 130-8644