

Read this sheet in full before using, and keep for later reference.

Find more product info online

BUFFERIN Luna i

Search

Fast, Effective Relief from Menstrual Pain and Headaches

<Pain reliever/fever reducer>

BUFFERIN Luna i

Designated Class 2 OTC drugs

Some BUFFERIN brand products contain different active ingredients. The pain relieving/fever reducing ingredients in this product are **ibuprofen** and **acetaminophen**. When consulting a doctor, dentist, pharmacist or registered seller about this product, specify that the product you are asking about contains ibuprofen and acetaminophen.

Quick and Effective

- Formulated with **ibuprofen** and **acetaminophen**
- Quick Melt capsules dissolve rapidly*

Contains no drowsiness-inducing ingredients

Small, easy-to-swallow tablets

Easy on the stomach



*Process patent 4065902, granulated pellets, tablets, and manufacturing methods of granulated pellets

Warnings

Do not

(Failure to comply with these instructions increases the likelihood of current symptoms worsening, side effects occurring, and other potentially serious risks)

1. Do not use this product if you

- (1) Have ever had an allergic reaction to this product or any of its ingredients
- (2) Have ever had an asthma attack after using this product, any other pain reliever/fever reducer, or a cold medicine
- (3) Are under 15 years old
- (4) Are pregnant and expecting to give birth within 12 weeks

2. Do not take any of the following pharmaceuticals while using this product

Other pain relievers/fever reducers, cold medicines, sedatives

3. Do not consume alcohol before or after taking this product

4. Do not use this product continuously for extended periods

Seek advice

1. Ask a doctor, dentist, pharmacist or registered seller before use if you are

- (1) Currently receiving treatment from a doctor or dentist
- (2) Pregnant or suspect you may be pregnant
- (3) Breast-feeding
- (4) Elderly

- (5) Have ever shown allergic symptoms when taking a drug or medicine
- (6) Have been diagnosed with any of the following conditions: heart disease; kidney disease; liver disease; systemic erythematosus; mixed connective tissue disease
- (7) Have ever had any of the following illnesses: stomach or duodenal ulcer; ulcerative colitis; Crohn's disease

2. The following symptoms may be side effects of the product. If you experience any of the following symptoms when using the product, stop use immediately and bring this sheet to a doctor, dentist,* pharmacist or registered seller for consultation.

Affected area	Symptom
Skin	Rash or redness; itchiness; bruising
Digestive tract	Nausea or vomiting; loss of appetite; stomach discomfort; stomach pain; oral inflammation; heartburn; heaviness in the stomach; gastrointestinal bleeding; abdominal pain; diarrhea; bloody stool
Mental state / Nervous system	Dizziness
Circulatory system	Heart palpitations
Respiratory system	Shortness of breath
Other	Blurred vision; ringing in the ears; swelling; nosebleed; bleeding from the gums; bleeding that does not stop; bleeding; back pain; excessive drop in body temperature; lethargy

The following rare but serious conditions may also occur. If you experience any of the symptoms below, seek immediate medical attention from a doctor.

Condition	Symptoms
Shock (anaphylaxis)	The following symptoms occur shortly after use: itchy skin; hives; hoarse voice; sneezing; itchy throat; difficulty breathing; heart palpitations; feeling faint
Mucocutaneous ocular syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, acute generalized exanthematous pustulosis	Persistence or sudden worsening of: severe fever; bloodshot eyes; eye mucus; lip sores; sore throat; widespread rash or skin redness; small pimples (pustules) appear on reddened skin; overall lethargy; loss of appetite
Impaired liver function	Fever; itchiness; rash; jaundice (skin and whites of the eyes appear yellowish); brown urine; overall lethargy; loss of appetite
Kidney disorder	Fever; rash; reduced urine output; swelling all over the body; overall lethargy, arthralgia (joint pain); diarrhea
Aseptic meningitis	Severe headache accompanied by neck stiffness; fever; nausea or vomiting (these symptoms are especially common in patients receiving care for systemic erythematosus or mixed connective-tissue disease)

Interstitial pneumonia	Sudden onset or persistence of: shortness of breath or difficulty breathing after mild exertion, such as when climbing stairs; dry coughing; fever
Asthma	Wheezing or gasping for breath, difficulty breathing
Aplastic anemia	Bruising; nosebleed; bleeding from the gums; fever; skin and mucous membranes appear pale; fatigue; heart palpitations; shortness of breath; feeling sick and dizziness; bloody urine
Agranulocytosis	Sudden onset of fever, chills or sore throat

- The following symptom may occur during use. If this symptom persists or worsens, stop use and bring this sheet to a doctor, pharmacist or registered seller for consultation.** Constipation
- If symptoms do not improve after 5 or 6 doses, stop use and bring this sheet to a doctor, dentist, pharmacist or registered seller for consultation.**

Effects

- Pain relieving effect on menstrual pain, headache, back pain, pain due to stiff shoulders, muscle pain, joint pain, pain from bruising, pain from bone fractures, pain from sprains, toothache, pain after tooth extraction, neuralgia, earache, pain from trauma and sore throat.
- Reducing effect on chills and fevers.

Directions and Dosage

Avoid taking on an empty stomach and wait at least four hours between doses.

Take the following amount with lukewarm or cold water.

Age	Adults (15 years and up)	Children under 15 years
Amount per 1 dose	2 tablets	Do not use
Doses per day	No more than 3	✘

Usage and Dosage Warnings

- Use only as directed.
- To remove the tablets from the package: As shown in the illustration on the right, press firmly with a fingertip on the protruding bubble in the clear plastic sheet that contains the tablet, pushing the tablet through the foil backing and out (swallowing the product without first removing it from the package can lead to injury, such as damage to the esophageal membrane).



Ingredients (per 2 tablets)

Active ingredient	Content	Purpose
Ibuprofen	130 mg	Reduces pain and fever
Acetaminophen	130 mg	
Anhydrous caffeine	80 mg	Boosts pain relief effect, eases headache

Dehydrated aluminum hydroxide gel	70 mg	Protects the gastric mucosa, helps prevent stomach problems
-----------------------------------	-------	---

Inactive ingredients: hydroxypropylcellulose, D-mannitol, lactic acid, magnesium stearate, polyvinyl alcohol (partially saponified), talc, titanium oxide and soy lecithin.

Storage and Handling

- (1) Store out of direct sunlight in a cool, dry place.
- (2) Keep out of reach of children.
- (3) Do not transfer to another container or package (may lead to incorrect use or affect product quality).
- (4) Do not use after the use-by date indicated on the package.
- (5) Avoid damaging the foil protecting tablets you are not planning to use immediately, as exposure can affect product quality.

Inquiries

Please direct inquiries to the store where you purchased the product or the service center below.

Lion Corporation Consumer Service Center [0120-813-752](tel:0120-813-752)

Hours available: 9:00–17:00 (excluding Saturdays, Sundays and holidays)

Lion Corporation

3-7, Honjo 1-chome, Sumida-ku, Tokyo 130-8644