

Read this sheet in full before using, and keep for later reference.

Find more product info online

BUFFERIN Luna J Search

Relief from Headaches and Menstrual Pain, OK for Teens and Children 7 and Up

<Pain reliever/fever reducer>

# BUFFERIN Luna J

Class 2 OTC drugs

Some BUFFERIN brand products contain different active ingredients. The pain relieving/fever reducing ingredient in this product is **acetaminophen**. When consulting a doctor, dentist, pharmacist or registered seller about this product, specify that the product you are asking about contains acetaminophen.

OK for use by children and teenagers with dosage adjusted by age!

Contains no drowsiness-inducing ingredients, so you can give your all at school.

Chewable tablet can be taken without water (non-bitter fruit flavor).

## Warnings

This product can be used by teenagers and children (age 7 and up). The warnings below are the general warnings for pain relievers/fever reducers.

### Do not

(Failure to comply with these instructions increases the likelihood of current symptoms worsening, side effects occurring, and other potentially serious risks)

#### 1. Do not use this product if you

- (1) Have ever had an allergic reaction to this product or any of its ingredients
- (2) Have ever had an asthma attack after using this product, any other pain reliever/fever reducer, or a cold medicine

#### 2. Do not take any of the following pharmaceuticals while using this product

Other pain relievers/fever reducers, cold medicines, sedatives

#### 3. Do not consume alcohol before or after taking this product

#### 4. Do not use this product continuously for extended periods

### Seek advice

#### 1. Ask a doctor, dentist, pharmacist or registered seller before use if you are

- (1) Currently receiving treatment from a doctor or dentist
- (2) Pregnant or suspect you may be pregnant
- (3) Elderly

- (4) Have ever shown allergic symptoms when taking a drug or medicine
- (5) Have been diagnosed with any of the following conditions: heart disease; kidney disease; liver disease; stomach or duodenal ulcer

**2. The following symptoms may be side effects of the product. If you experience any of the following symptoms when using the product, stop use immediately and bring this sheet to a doctor, pharmacist or registered seller for consultation.**

Affected area	Symptom
Skin	Rash or redness; itchiness
Digestive tract	Nausea or vomiting; loss of appetite
Mental state / Nervous system	Dizziness
Other	Excessive drop in body temperature

The following rare but serious conditions may also occur. If you experience any of the symptoms below, seek immediate medical attention from a doctor.

Condition	Symptoms
Shock (anaphylaxis)	The following symptoms occur shortly after use: itchy skin; hives; hoarse voice; sneezing; itchy throat; difficulty breathing; heart palpitations; feeling faint
Mucocutaneous ocular syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, acute generalized exanthematous pustulosis	Persistence or sudden worsening of: severe fever; bloodshot eyes; eye mucus; lip sores; sore throat; widespread rash or skin redness; small pimples (pustules) appear on reddened skin; overall lethargy; loss of appetite
Impaired liver function	Fever; itchiness; rash; jaundice (skin and whites of the eyes appear yellowish); brown urine; overall lethargy; loss of appetite
Kidney disorder	Fever; rash; reduced urine output; swelling all over the body; overall lethargy, arthralgia (joint pain); diarrhea
Interstitial pneumonia	Sudden onset or persistence of: shortness of breath or difficulty breathing after mild exertion, such as when climbing stairs; dry coughing; fever
Asthma	Wheezing or gasping for breath, difficulty breathing

**3. If symptoms do not improve after 5 or 6 doses, stop use and bring this sheet to a doctor, dentist, pharmacist or registered seller for consultation.**

### Effects

- (1) Pain relieving effect on menstrual pain, headache, back pain, pain due to stiff shoulders, muscle pain, joint pain, pain from bruising, pain from bone fractures, pain from sprains, toothache, pain after tooth extraction, neuralgia, earache, pain from trauma and sore throat.
- (2) Reducing effect on chills and fevers.

## Directions and Dosage

Avoid taking on an empty stomach. Chew or dissolve in the mouth. Wait at least four hours between doses.

Age	<u>15 and up</u>	11 to 14	7 to 10	Under 7
Amount per 1 dose	<u>3 tablets</u>	2 tablets	1 tablet	Do not use
Doses per day	No more than 3			<b>X</b>

### Usage and Dosage Warnings

- (1) For administration to children under the supervision and guidance of a guardian or caretaker. When given to children in advance for later use when away from direct supervision, provide thorough instruction on correct usage and dosage ahead of time.
- (2) To remove the tablets from the package: As shown in the illustration on the right, press firmly with a fingertip on the protruding bubble in the clear plastic sheet that contains the tablet, pushing the tablet through the foil backing and out (swallowing the product without first removing it from the package can lead to injury, such as damage to the esophageal membrane).
- (3) Use only as directed.



## Ingredients (per 1 tablets)

Active ingredient	Content	Purpose
Acetaminophen	100 mg	Reduces pain and fever

Inactive ingredients: ethyl cellulose, sodium lauryl sulfate, cetanol, triacetin, synthetic hydrotalcite, D-mannitol, cornstarch, hydroxypropylcellulose, sucralose, l-menthol, flavoring, dextrin, vanillin, and magnesium stearate.

### Storage and Handling

- (1) Store out of direct sunlight in a cool, dry place.
- (2) Keep out of reach of children.
- (3) Do not transfer to another container or package (may lead to incorrect use or affect product quality).
- (4) Do not use after the use-by date indicated on the package.
- (5) Avoid damaging the foil protecting tablets you are not planning to use immediately, as exposure can affect product quality.

### Inquiries

Please direct inquiries to the store where you purchased the product or the service center below.

Lion Corporation Consumer Service Center [0120-813-752](tel:0120-813-752)

Hours available: 9:00–17:00 (excluding Saturdays, Sundays and holidays)

## Lion Corporation

3-7, Honjo 1-chome, Sumida-ku, Tokyo 130-8644